



Cette présentation a été effectuée le 30 novembre 2004, au cours de la journée « Obstétrique et santé publique : élargir les perspectives sur les réalités de la naissance » dans le cadre des Journées annuelles de santé publique (JASP) 2004. L'ensemble des présentations est disponible sur le site Web des JASP, à l'adresse <http://www.inspq.qc.ca/jasp/archives/>.









Support émotionnel



Support



Physique









Réduction de 50%
des taux de
césarienne

40% moins
d'utilisation
d'ocytocine

Accouchements
25% plus courts

Diminution
de 30% d'utilisation
d'analgésiques

Réduction de
40% de
l'utilisation de
forceps

Réduction de
60% des
demandes de
péridurale

Kennell J, Klaus M, McGrath S, Robertson S, Jinkley C.
Medical Intervention : The Effect of Social Support During Labor, 1988



Hodnett ED, Gates S, Hofmeyr G J, Sakala C.
Continuous Support for Women During Childbirth, 2003



Hodnett ED, Lowe NK, Hannah ME, Willan AR, Stevens B et al.
Effectiveness of Nurses as Providers of Birth Labor Support in North American Hospitals, 2002



Hodnett ED, Lowe NK, Hannah ME, Willan AR, Stevens B et al.
Effectiveness of Nurses as Providers of Birth Labor Support in North American Hospitals, 2002

